

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/123437/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Allen, Lowri A, Pyart, Rhodri D, Holmes, Jennifer, Donovan, Kieron L, Anderson, Richard A and Phillips, Aled O ORCID: <https://orcid.org/0000-0001-9744-7113> 2019. Cardiovascular and renal outcomes following percutaneous coronary intervention in a population with renal disease: a case-control study. QJM: An International Journal of Medicine 112 (9) , pp. 669-674. 10.1093/qjmed/hcz130 file

Publishers page: <http://dx.doi.org/10.1093/qjmed/hcz130>  
<<http://dx.doi.org/10.1093/qjmed/hcz130>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Cardiovascular and renal outcomes following percutaneous coronary intervention in a population  
with renal disease: a case-control study

Lowri A Allen, MBChB<sup>1</sup>, Rhodri D Pyart, MBBCh<sup>1</sup>, Jennifer Holmes<sup>2</sup>, MSc<sup>1</sup>, Kieron L Donovan, MD<sup>1</sup>,  
Richard A Anderson, MD<sup>3</sup>, Aled O Phillips, MD<sup>1</sup>.

<sup>1</sup>Institute of Nephrology, Cardiff University School of Medicine, Cardiff, UK.

<sup>2</sup>Welsh Renal Clinical Network, Cwm Taf University Health Board, Abercynon, UK.

<sup>3</sup>Department of Cardiology, Cardiff University School of Medicine, Cardiff, UK.

Corresponding author:

Prof A O Phillips

Institute of Nephrology

Cardiff University School of Medicine

University Hospital of Wales

Heath Park, Cardiff

CF14 4XN

Tel: 02920 748467

Fax: 02920 748470

E-mail: phillipsao@cf.ac.uk

Word count:

Abstract: 250

Main body: 2379 (excluding abstract, references, acknowledgements and figures)

**Abstract:**

**Background:** Patients with renal disease are less likely to undergo percutaneous coronary intervention (PCI) due to concerns about poor outcomes.

**Aim:** We describe outcomes following PCI in individuals with chronic kidney disease (CKD), as compared with matched controls with comparable CKD who did not undergo PCI. We also identified factors predictive of poor outcomes following PCI amongst patients with CKD.

**Design:** Retrospective observational case-control study.

**Methods:** Cases were individuals with CKD (stages 1-5) undergoing PCI between 2008-2014. Controls were age, gender and creatinine matched individuals not requiring PCI. We compared mortality between groups using Kaplan Meier curves and Cox regression modelling. We assessed changes in serum creatinine using Wilcoxon Rank testing. We explored the relationship between biochemical and haematological measures (baseline creatinine, calcium, phosphate, calcium-phosphate product, parathyroid hormone, white cell count, haemoglobin, platelet count, c-reactive protein, total cholesterol) and post-PCI mortality, using logistic regression.

**Results:** We identified 144 cases and 144 controls. Mortality was significantly lower amongst cases compared with controls (hazard ratio 0.46 (95% confidence intervals 0.31,0.69)). PCI did not result in a significant change in renal function ( $p=0.52$ ). Amongst cases, serum creatinine and calcium-phosphate product were predictors of mortality following PCI.

**Conclusion:** Cases undergoing PCI had lower mortality, and PCI was not associated with accelerated CKD progression. On this data, PCI should not be deferred as a treatment option in patients with

- 1 CKD. Serum creatinine and calcium-phosphate product predict mortality following PCI in this cohort,
- 2 and may be useful in risk-stratifying patients with CKD being considered for PCI.
- 3

## **Background:**

Cardiovascular disease is one of the leading causes of mortality in patients with chronic kidney disease (CKD)<sup>1-4</sup>. Cardiovascular mortality is 10-20 times more common in patients with CKD Stage 5 who are on dialysis, compared with the background population<sup>2</sup>. Furthermore, there is evidence to suggest that cardiovascular disease typically occurs at an earlier age, and follows a more accelerated course in patients with CKD, as compared with the background population<sup>5,6</sup>. Unlike in the background population, cardiovascular disease is not typically the result of progressive atherosclerosis, but instead is due to dysregulation of calcium and phosphate homeostasis, and the development of medial vascular calcification<sup>7-9</sup>.

There is no consensus regarding how best to manage cardiovascular disease in patients with CKD. Most trials of treatments for cardiovascular disease exclude patients with renal disease, and as such the evidence base underpinning the management of these patients' cardiovascular disease is sparse<sup>10,11</sup>. In practice, patients with CKD are less likely to receive medications such as Aspirin, ACE inhibitors and Beta blockers, due to concerns about side effects/toxicity and adverse effects on renal function<sup>10,12</sup>. In addition, patients with CKD are less likely to be offered angiography and or percutaneous coronary intervention (PCI)<sup>12</sup>. This is in part a reflection of the fact that renal impairment predicts poorer outcomes including mortality and procedural complications such as bleeding and in-stent stenosis following PCI<sup>12-15</sup>. Furthermore, many clinicians are concerned that the contrast load associated with angiography can lead to an accelerated loss of renal function and an earlier need for renal replacement therapy<sup>16,18</sup>.

Whilst many studies have explored outcomes following PCI for patients with CKD compared with patients with normal renal function, few have compared outcomes for patients with CKD undergoing PCI and those with comparable CKD not requiring PCI<sup>13,14,19,20</sup>. As such, it is unclear whether the poor

outcomes previously described amongst patients with renal disease undergoing PCI are a result of the PCI, or simply a marker of the high degree of morbidity and mortality associated with CKD itself.

This study aimed to describe mortality, cardiovascular and renal outcomes following PCI in patients with CKD, as compared with age, sex and creatinine matched controls who had not undergone PCI. We also explored the relationship between baseline biochemistry and haematology results and mortality following PCI. In particular, we explored whether serum measurements of calcium, phosphate and/or calcium-phosphate product are useful predictors of mortality amongst this cohort, as dysregulation of calcium and phosphate metabolism is thought to be the main reason for the medial vascular calcification that drives excess cardiovascular morbidity and mortality in patients with renal disease.

## **Methods:**

### *Study population:*

We conducted a retrospective case-control study. Cases were individuals who had undergone percutaneous coronary intervention between 1<sup>st</sup> January 2008 and 31<sup>st</sup> December 2014, and who were known to the nephrology team at the time of intervention. Cases were identified by cross-checking all individuals who underwent PCI in a tertiary cardiology centre (University Hospital of Wales) during the study period, with a database of patients receiving care under a nephrology service (Vital Data). Controls were individuals receiving care from nephrology services who had not undergone percutaneous coronary intervention. Controls were age, gender and creatinine matched. For individuals who underwent PCI, the indication for PCI, number of vessels treated, and location of lesions treated were documented. Data regarding left ventricular function was not available in this retrospective study. To minimise the risk of contrast induced nephropathy, the lowest volume of low osmolar contrast agent was used for all individuals who underwent PCI, in line with current guidelines<sup>21</sup>. We compared outcomes for cases and controls.

### *Outcomes:*

Cases were followed up from the time of PCI to the end of the data collection period (1<sup>st</sup> July 2016) or until the time of death. Controls were followed up from the date of the blood test upon which age and creatinine-based matching was based until the end of the data collection period (1<sup>st</sup> July 2016) or until the time of death.

The primary outcome was mortality at the end of follow up. We also collected data regarding changes in serum creatinine following PCI, to ascertain whether PCI was associated with acceleration in the progression of CKD in the long term. We identified cases of contrast induced nephropathy using the recognised definition of a relative (>25%) or absolute ( $\geq 44 \mu\text{mol/l}$ ) increase in serum

1 creatinine from the baseline value within the first 48-72hours following percutaneous coronary  
2 intervention<sup>22</sup>.

3

4 We also collected data on cardiovascular outcomes amongst cases. Specifically, we documented  
5 whether they had experienced recurrent angina; an acute coronary syndrome; symptomatic heart  
6 failure, or if they required further angiography/PCI prior to the end of the follow up period.

7



*Secondary analysis:*

We collected data regarding baseline creatinine, calcium, phosphate, calcium-phosphate product, parathyroid hormone, white cell count, haemoglobin, platelet count, c-reactive protein, and total cholesterol for all cases. Baseline measures were defined as the mean value of all results during the 3 months preceding PCI. We explored the relationship between these variables and mortality.

*Statistical methods:*

Survival time was compared between cases and controls using Kaplan Meier survival curves. Cox regression modelling was subsequently used to obtain an adjusted hazard ratios (aHR) with 95% confidence intervals. Data were right censored at the end date of the follow up period for patients who remained alive.

For cases, we used a Wilcoxon rank test to compare serum creatinine at baseline with serum creatinine at the end of the follow up period (due to the non-parametric distribution of the data).

The proportion of cases experiencing adverse cardiovascular outcomes were described.

Logistic regression modelling was used to explore the relationship between the baseline haematology and biochemistry results of cases and mortality following PCI. Estimates of effects between cases and controls are reported as crude odds ratios (ORs) and adjusted ORs (aORs) alongside 95% confidence intervals. Adjusted odd ratios were derived after adjusting for age, gender and baseline creatinine as appropriate.

Ethics and Information Governance: Patient data and outcomes were retrieved from internal databases. This work represents an audit of outcomes in our department, and as such ethical approval was not required.

- 1 Data analyses were carried out using IBM SPSS Statistics version 22 and Stata version 14.

## **Results:**

We identified 144 patients who underwent PCI between 1<sup>st</sup> January 2008 and 31<sup>st</sup> December 2014, who were known to the nephrology team at the time of intervention. 144 age, gender and creatinine matched controls were identified. The mean follow-up period was 42.2 months for cases (range 1-100 months) and 36.0 months for controls (range 0-87 months.)

Baseline characteristics are described in table 1. Characteristics were similar for cases and controls, indicating appropriate matching of cases to controls. Indications for percutaneous coronary intervention amongst cases are described in table 2a. The commonest indication for PCI was acute coronary syndrome, followed by angina. A small proportion of cases underwent elective angiography and percutaneous coronary intervention in the absence of any symptoms or signs of coronary ischaemia, as part of the routine work up for renal transplantation and subsequent demonstration of ischaemia.

The cases included 106 patients treated for single vessel coronary artery disease, 35 patients with 2 vessel disease and 1 with 3 vessel disease (table 2b). Lesions treated were located in the right coronary artery lesions in 60 cases, the left circumflex in 17 cases and the left anterior descending artery in 51 cases, and the left main stem vessels in 4 cases (table 2c). The remainder of treated lesions related to side branch disease involving either large diagonal vessel or obtuse marginals.

PCI was associated with lower mortality compared to the non-PCI matched control group. Mortality was high in both groups (28% at the end of follow up in cases, and 55% at the end of follow up amongst controls). Time to mortality was lower amongst the control group, and this was statistically significant (hazard ratio 0.46 (0.31, 0.69)) (figure 1.)

1 PCI did not lead to long-term deterioration in renal function. There was no statistically significant  
2 change in creatinine amongst cases following percutaneous coronary intervention (mean creatinine  
3 prior to intervention = 198 micromol/L, mean creatinine at the end of follow up =196 micromol/L,  
4  $p=0.52$ .)

5  
6 Data regarding contrast induced nephropathy was available for 88 (61%) of cases, of whom 9  
7 (10.2%) met the criteria for a diagnosis of contrast induced nephropathy.

8  
9 Cardiac outcomes for cases are described in table 2b. Most patients had a favourable outcome.  
10 82% remained free of symptomatic cardiovascular disease at the end of the follow up period. The  
11 commonest adverse cardiovascular outcome was a recurrence of angina, affecting 19% of patients.

12  
13 Univariate analysis revealed increased age, calcium, phosphate, and calcium-phosphate product to  
14 be associated with a statistically significant increased risk of mortality amongst cases. Increased  
15 haemoglobin was associated with a lower risk of mortality (table 3).

16  
17 After adjusting for age and gender, an increased creatinine continued to be associated with  
18 increased mortality after PCI ( $p=0.003$ .) After adjusting for age, gender and creatinine, serum  
19 phosphate and haemoglobin were no longer significantly associated with mortality. However,  
20 calcium-phosphate product was still significantly associated with an increased mortality ( $p=0.04$ )  
21 (table 3).

## **Discussion**

Our results confirm that mortality is high amongst patients with CKD, as would be expected from the results of previously published studies<sup>23,24</sup>. Whilst studies have previously demonstrated high mortality amongst patients with CKD undergoing PCI as compared with individuals with normal renal function undergoing PCI, it is unclear whether the excess mortality is a result of PCI, or whether it reflects the high levels of mortality associated with renal disease itself. In order to elucidate this, it is necessary to compare outcomes between individuals with renal disease undergoing PCI, and individuals with comparable CKD not requiring PCI. Our study used a case-control approach to compare outcomes between individuals with renal disease undergoing PCI and creatinine-matched controls who did not undergo PCI.

The first key finding from our study is that patients with renal disease who underwent PCI had lower mortality than creatinine-matched controls. This data would suggest that PCI may protect against short term mortality in a cohort of patients with renal disease. This is in-keeping with the findings of a previously published systematic review which concluded that early angiography and/or PCI conferred a survival benefit, at least in those with early-moderate CKD<sup>25</sup>. We therefore suggest that the excess of mortality following PCI previously documented amongst patients with renal disease is likely to reflect the poor prognosis associated with having renal disease in itself, rather than being directly related to the PCI procedure.

In addition to the perceived high rates of mortality amongst CKD patients undergoing PCI, many clinicians have expressed concern that PCI may lead to an accelerated loss of renal function amongst patients with CKD as a result of the contrast burden associated with the procedure<sup>16,18</sup>. The concern that PCI may hasten time to end stage renal failure and the need for renal replacement therapy has contributed to the reluctance to offer PCI to many patients with renal disease<sup>12</sup>. 10.2% of cases developed contrast induced nephropathy, which is comparable with the incidence of contrast

1 induced nephropathy in previously published studies<sup>26,27</sup>. Importantly we did not observe an  
2 accelerated rate of decline in renal function in the long-term following PCI amongst our cases.  
3 Whilst one of the limitations of this study is the relatively short follow up period (mean 42.24  
4 months, range 1-100 months), deteriorating renal function as a direct result of PCI is usually due to  
5 an acute effect of contrast and associated acute kidney injury, and would have been apparent during  
6 this period of follow up<sup>16,18</sup>.

7  
8 Since our data suggests that PCI confers a survival advantage in this group, and is not associated with  
9 an accelerated decline in renal function amongst patients with CKD, we suggest that having CKD in  
10 itself should not be considered a contraindication to PCI. Furthermore, since patients with renal  
11 disease have high mortality rates independently of the need for coronary revascularisation, much of  
12 the focus of care for patients with progressive renal disease is on trying to maximise quality of life  
13 and minimise distressing symptoms<sup>23,24</sup>. Amongst our cohort, 81% did not experience recurrent  
14 angina during the follow up period, thereby suggesting PCI had a positive impact on their symptom  
15 profile, at least in the short term. We therefore believe that PCI is a treatment that has the potential  
16 to dramatically improve quality of life in patients with renal disease, without accelerating the  
17 progression of their underlying renal disease.

18  
19 We recognise that our case mix includes patients with a broad spectrum of renal disease. We  
20 therefore sought to identify factors associated with an increased risk of mortality amongst our cases.  
21 As expected, we found that increased creatinine was associated with a higher mortality post  
22 procedure, independently of age and gender. Similarly, a high serum phosphate and low serum  
23 haemoglobin, both of which are markers of severe renal disease, were significantly associated with  
24 higher mortality. After correcting for creatinine, neither remained significantly associated with  
25 mortality, illustrating the fact that they act as surrogate markers of the severity of renal disease in

1 this setting. These findings support our observation that the severity of renal disease is one of the  
2 main predictors of mortality amongst this cohort.

3  
4 The only biochemical marker that remained significantly associated with mortality, after adjusting  
5 for age, gender and creatinine, was the calcium-phosphate product. It is well recognised that in  
6 patients with CKD, the excess cardiovascular mortality is not due to the traditional model of  
7 atherosclerotic disease, but is instead a result of vascular calcification that occurs secondary to  
8 disruption of calcium and phosphate homeostasis<sup>7-9</sup>. Studies have previously demonstrated that  
9 calcium-phosphate product may be a useful tool in estimating the risk of mortality and  
10 cardiovascular disease in patients with CKD<sup>28-30</sup>. Our data suggest that calcium-phosphate product  
11 may also be useful in predicting mortality following PCI in this group. In addition, since calcium-  
12 phosphate product may be affected by several modifiable factors such as diet, the use of phosphate  
13 binders and effective vitamin D replacement, it is possible that efforts to lower calcium-phosphate  
14 product could lead to more favourable outcomes in this group. Further studies are required to  
15 explore this<sup>30</sup>.

16  
17 Although this study to our knowledge is the first to compare the outcome of PCI in CKD patients  
18 compared to an age matched control group, a limitation of this study is that the majority of cases  
19 underwent PCI to treat single vessel disease with the highest proportion of treated vessels being  
20 RCA. , We acknowledge therefore that we should be cautious in generalising the findings to  
21 individuals undergoing PCI for multi vessel disease who by definition have more severe coronary  
22 artery disease. ~~and are likely to have higher levels of exposure to contrast during the procedure.~~  
23 However, our findings remain relevant to clinical practice, as data from the British Cardiovascular  
24 Intervention Society Adult Interventional Procedure Audit 2016 demonstrate that nationally, the  
25 vast majority of PCI procedures are undertaken to treat single vessel disease (77.2% of all PCI)<sup>31</sup>.  
26 ~~Whilst in our study the commonest location of lesions treated was the right coronary artery,~~

1 nationally the commonest location for lesions treated by PCI is the left anterior descending artery<sup>21</sup>.

2 Further studies are however required to establish whether there is an association between the  
3 location of lesions treated and outcomes in patients with chronic kidney disease.

#### 5 **Summary**

6 We have demonstrated that PCI was associated with a reduction in mortality, and did not lead to an  
7 acceleration in the progression of CKD in a cohort of patients with CKD compared with creatinine  
8 matched controls who did not undergo percutaneous coronary intervention. We therefore suggest  
9 that CKD in itself should not be considered a contraindication to PCI. The decision to undertake PCI  
10 should be considered on an individual patient basis, and should take into account the severity of  
11 their CKD. Calcium-phosphate product may be a useful tool in identifying patients with renal disease  
12 who are at high risk of mortality following PCI.



1    **Acknowledgments:** Nil.

2    **Funding:** Nil.

3    **Conflict of interest:** None of the authors have any conflicts of interest or disclosures.

4    **Author contributions:** LAA was involved in collating the data for cases and was responsible for

5    undertaking statistical analysis and writing the manuscript. RDP was responsible for identifying

6    matched controls, collating the data for controls, and also provided statistics advice and support. JH

7    was responsible for identifying cases and extracting biochemistry and haematology data from

8    internal databases. KLD, RAA and AOP supervised the project, provided critical feedback and helped

9    shape the research, analysis and manuscript. All authors discussed the results and contributed to

10   the final manuscript.

## 1 References

- 2 1. Foley RN, Parfrey PS, Sarnak MJ. Epidemiology of cardiovascular disease in chronic renal  
3 disease. JASN. 1998;9;S16-23.
- 4 2. Sarnak MJ, Levey AS. Cardiovascular disease and chronic renal disease: A new paradigm.  
5 AJKD. 2000;35;S117-131
- 6 3. Collins AJ. Cardiovascular mortality in end-stage renal disease. Am J Med Sci. 2003;4;163-  
7 167.
- 8 4. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu C. Chronic kidney disease and the risks of  
9 death, cardiovascular events, and hospitalization. N Eng J Med. 2004;351;1296-1305.
- 10 5. Schifflin EL, Lipman ML, Mann JFE. Chronic kidney disease: effects on the cardiovascular  
11 disease. Circulation. 2007;116;85-97.
- 12 6. Baigent C, Burbury K, Wheeler D. Premature cardiovascular disease in chronic renal failure.  
13 Lancet. 2000;356;147-152.
- 14 7. London GM, Guerin AP, Marchais S, Metivier F, Pannier B, Adda H: Arterial medial  
15 calcification in endstage renal disease: Impact on all-cause and cardiovascular mortality.  
16 Nephrol Dial Transplant. 2003;18;1731–1740.
- 17 8. Blacher J, Guerin AP, Pannier B, Marchais SJ, London GM: Arterial calcifications, arterial  
18 stiffness, and cardiovascular risk in end-stage renal disease. Hypertension. 2001;38; 938–  
19 942.
- 20 9. Wilson PW, Kauppila LI, O'Donnell CJ, Kiel DP, Hannan M, Polak JM, Cupples LA: Abdominal  
21 aortic calcific deposits are an important predictor of vascular morbidity and mortality.  
22 Circulation. 2001;103;1529–1534.
- 23 10. Ezekowitz J, McAlister FA, Humphries KH, Norris CM, Tonelli M, Ghali WA, Knudston ML,  
24 APPROACH Investigators. The association among renal insufficiency, pharmacotherapy, and  
25 outcomes in 6,427 patients with heart failure and coronary artery disease. JACC. 2004; 44;  
26 1587-1592.

11. Coca SG, Krumholz HM, Garg AX, Parikh CR. Underrepresentation of renal disease in randomized controlled trials of cardiovascular disease. *JAMA*. 2006;296:1377-84.
12. Masoudi FA, Plomondon ME, Magid DJ, Sales A, Rumsfield JS. Renal insufficiency and mortality from acute coronary syndromes. *Am Heart J*. 2004; 147:623-629.
13. Best PJM, Lennon R, Ting HH, Bell MR, Rihal CS, Holmes DR, Berger PB. The impact of renal insufficiency on clinical outcomes in patients undergoing percutaneous coronary interventions. *JACC*. 2002;39:1113-1119
14. Sadeghi HM, Stone GW, Grines CL, Mehran R, Dixon SR, Lansky AJ, Fahy M, Cox DA, Garcia E, Tcheng JE, Griffin JJ, Stuckey TD, Turco M, Carroll JD. Impact of renal insufficiency in patients undergoing primary angioplasty for acute myocardial infarction. *Circulation*. 2003;108; 2769-2775.
15. Medi C, Montalescot G, Budaj A, Fox LA, Lopez-Sendon J, FitzGerald G, Brieger DB; GRACE Investigations. Reperfusion in patients with renal dysfunction after presentation with ST-segment elevation or left bundle branch block: GRACE (Global Registry of Acute Coronary Events). *JACC Cardiovasc Interv*. 2009;2;26-33.
16. Widimsky P, Rychlik I. Renal disease and acute coronary syndrome. *Heart*. 2010;96;86-92.
17. Finn WF. The clinical and renal consequences of contrast-induced nephropathy. *Nephrol Dial Transplant*. 2006; 21: 2-10.
18. McCullough P. Outcomes of contrast-induced nephropathy. Experience in patients undergoing cardiovascular intervention. *Coronary artery dis*. 2006; 67:334-343.
19. Latif F, Kleiman NS, Cohen DJ, Pencina MJ, Yen CH, Cutlip DW, Moliterno DJ, Nassif D, Lopez JL, Saucedo JF, EVENT Investigators. In-hospital and 1-year outcomes among percutaneous coronary intervention patients with chronic kidney disease in the era of drug-eluting stents: a report from the EVENT registry. *JACC: Cardiovascular interventions*. 2009;2:37-45.

20. Gruberg L, Dangas G, Mehran R, Mintz GS, Kent KM, Pichard AD, Satler LF, Lansky AJ, Stone GW, Leon MB. Clinical outcome following percutaneous coronary intervention in patients with chronic renal failure. *Catheterisation and cardiovascular interventions*. 2002;55:66-72.
21. Neumann FJ, Sousa-Uva M, Ahlsson A, Alfonso F, Bannin AP, Bendetto U, Byrne RA, Collet JP, Falk V, Juni SJHP, Kastrati A, Holler A, Kristensen SD, Niebauer J, Ritcher DJ, Seferovic PM, Sibbing D, Stefanini GG, Windecker S, Yadav R, Zembala MO. ESC/EACTS Guidelines on Myocardial Revascularisation. *Eur Heart J* 2019;40:87-165.
22. Mehran R, Aymong ED, Nikolsky E, Lasic Z, Iakovou I, Fahy M, Mintz GS, Lansky AJ, Moses JW, Stone GW, Leon MB, Dangas G. A simple risk score for prediction of contrast-induced nephropathy after percutaneous coronary intervention: development and initial validation. *J Am Coll Cardiol* 2004;44:1393 – 1399.
23. Tonelli M, Wiebe N, Culleton B, House A, Rabbat C, Fok M, McAlister F, Garg AX. Chronic kidney disease and mortality risk: a systematic review. *JASN*. 2006;17:2034-2047.
24. Keith DS, Nichols GA, Gullion CM, Brown JB, Smith DH. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organisation. *Arch Intern Med*. 2004;164:659-663.
25. Shaw C, Nitsch D, Lee J, Fogarty D, Sharpe CC. Impact of an early invasive strategy versus conservative for unstable angina and Non-ST elevation acute coronary syndrome in patients with chronic kidney disease: a systematic review. *PLoS One*, 2016;11:p. e0153478.
26. Mehran R, Aymong ED, Nikolsky E, Lasic Z, Iakovour I, Fahy M, Mintz GS, Lansky AJ, Moses JW, Stone GW, Leon MB, Dangas G. A simple risk score for prediction of contrast-induced nephropathy after percutaneous coronary intervention. *JACC*.2004;7:1393-1399
27. McCullough PA, Adam A, Becker CR, Davidson C, Lameire N, Stacul F, Tumlin J, CIN Consensus Working Panel. *The American Journal of Cardiology*. 2006;6: 5-13.

- 1 28. Block GA, Hulbert-Shearon TE, Levin NH, Port FK. Association of serum phosphorus and  
2 calcium x phosphate product with mortality risk in chronic haemodialysis patients: a national  
3 study. *Am J Kidney Dis.* 1998;31:607-17.
- 4 29. Ganesh SK, Stack AG, Levin NW, Hulbert-Shearon TE, Port FK. Association of elevated serum  
5 PO(4), Ca x PO(4) product, and parathyroid hormone with cardiac mortality risk in  
6 haemodialysis patients. *J Am Soc Nephrol.* 2001;12:2131-8.
- 7 30. Cozzolino M, Dusso AS, Slatopolsky E. Role of calcium-phosphate product and bone-  
8 associated proteins on vascular calcification in renal failure. *JASN.* 2001;12:2511-2516.
- 9 31. Ludman PF on behalf of the British Cardiovascular Intervention Society. BCIS Audit Returns;  
10 Adult Interventional Procedures. 2016. Available from [http://www.bcis.org.uk/wp-](http://www.bcis.org.uk/wp-content/uploads/2018/03/BCIS-Audit-2016-data-ALL-excluding-TAVI-08-03-2018-for-web.pdf)  
11 [content/uploads/2018/03/BCIS-Audit-2016-data-ALL-excluding-TAVI-08-03-2018-for-](http://www.bcis.org.uk/wp-content/uploads/2018/03/BCIS-Audit-2016-data-ALL-excluding-TAVI-08-03-2018-for-web.pdf)  
12 [web.pdf](http://www.bcis.org.uk/wp-content/uploads/2018/03/BCIS-Audit-2016-data-ALL-excluding-TAVI-08-03-2018-for-web.pdf). Accessed April 2019.
- 13

	Cases	Controls
Age (years)	69 (62.0, 77.0)	69 (61.7, 77.4)
Creatinine (micromol/L)	142 (103, 206)	142 (102, 209)
Gender	103 (72%)	103 (72%)
Diabetes	41 (28.4%)	45 (31.3%)

- 1 **Table 1:** Baseline characteristics of cases and controls. Values are median, interquartile range (as
- 2 data was positively skewed). Gender is described as the number and % of males in each group. The
- 3 number and % of individuals with co-existing diabetes are also described.

1

	Proportion of cases
Acute coronary syndrome	86 (60%)
Angina	42 (29%)
Investigation of heart failure	3 (2%)
Pre renal transplant	6 (4%)
Ventricular Tachycardia of unknown origin	2 (1%)
Unknown	5 (4%)

2 **Table 2a:** Indication for percutaneous coronary intervention.

	Proportion of cases
Single vessel coronary artery disease treated	106 (74%)
Two vessel disease treated	35 (24%)
Triple vessel disease treated	1 (1%)

3 **Table 2b:** Number of vessels treated.

	Proportion of lesions
Right coronary artery	60 (34%)
Left circumflex artery	17 (10%)
Left anterior descending artery	51 (29%)
Left main stem vessels	4 (2%)
Side branch disease (large diagonal vessel or obtuse marginals)	45 (25%)

4 **Table 2c:** Location of lesions treated

5

6

7

8

	Proportion of cases
Death	40 (27.8%)
Recurrent angina	27 (18.8%)
Recurrent ACS	14 (9.7%)
Heart failure	18 (12.5%)
Repeat angiogram needed	26 (18.1%)
Further PCI	23 (16%)

1 **Table 3:** Cardiovascular outcomes following percutaneous coronary intervention.




	Odds Ratio ± 95% CI	p value	Adjusted Odds Ratio ± 95% CI	p value
Age	1.60 (1.16, 2.23)	0.01		
Gender	0.94 (0.41, 2.11)	0.87		
Creatinine	1.00 (1.00, 1.01)	0.01	1.00 (1.00, 1.01)*	0.003
Calcium	0.25 (0.01, 4.92)	0.81	0.78 (0.03, 20.69)**	0.877
Phosphate	5.57 (1.56, 19.89)	0.008	6.43 (0.97, 42.64)**	0.054
Calcium Phosphate product	2.32 (1.24, 4.35)	0.008	2.34 (1.03, 5.34)**	0.042
PTH	1.04 (0.99, 1.09)	0.15	1.04 (0.99, 1.10)**	0.146
Hb	0.97 (0.94, 0.99)	0.005	0.98 (0.95, 1.01)**	0.220
WCC	1.05 (0.90, 1.24)	0.53	1.08 (0.91, 1.28)**	0.399
Platelets	1.00 (0.96, 1.00)	0.89	1.00 (1.00, 1.01)**	0.485
CRP	1.00 (0.99, 1.02)	0.62	1.00 (0.99, 1.02)**	0.784
Total Cholesterol	1.17 (0.81, 1.70)	0.40	1.27 (0.84, 1.92)**	0.267

2 **Table 3:** Relationship between baseline biochemistry and haematology results and mortality.

3 Results are crude and adjusted odds ratios with 95% confidence intervals and p values derived from  
4 logistic regression models.

5 \*Adjusted for age and gender

6 \*\* Adjusted for age, gender, creatinine



1

2 **Figure 1:** Kaplan Meier Curves demonstrating time to mortality (in days) amongst cases and

3 controls.